

# QIAEX II® Handbook

For DNA extraction from agarose and polyacrylamide gels and for desalting and concentrating DNA from solutions



# QIAGEN Sample and Assay Technologies

QIAGEN is the leading provider of innovative sample and assay technologies, enabling the isolation and detection of contents of any biological sample. Our advanced, high-quality products and services ensure success from sample to result.

## **QIAGEN sets standards in:**

- Purification of DNA, RNA, and proteins
- Nucleic acid and protein assays
- microRNA research and RNAi
- Automation of sample and assay technologies

Our mission is to enable you to achieve outstanding success and breakthroughs. For more information, visit [www.qiagen.com](http://www.qiagen.com).

# Contents

<b>Kit Contents</b>	<b>4</b>
<b>Storage</b>	<b>4</b>
<b>Product Use Limitations</b>	<b>4</b>
<b>Product Warranty and Satisfaction Guarantee</b>	<b>5</b>
<b>Technical Assistance</b>	<b>5</b>
<b>Safety Information</b>	<b>6</b>
<b>Product Specifications</b>	<b>6</b>
<b>Quality Control</b>	<b>6</b>
<b>Introduction</b>	<b>7</b>
Principle	7
<b>Protocols</b>	
■ Extraction of 40 bp to 50 kb of DNA Fragments from Agarose Gels	12
■ Extraction of DNA Fragments from Polyacrylamide Gels	15
■ Desalting and Concentrating DNA Solutions	17
<b>Troubleshooting Guide</b>	<b>19</b>
<b>References</b>	<b>21</b>
<b>Ordering Information</b>	<b>22</b>

## Kit Contents

<b>QIAEX II Gel Extraction Kit</b>	<b>(150)</b>	<b>(500)</b>
<b>Catalog no.</b>	<b>20021</b>	<b>20051</b>
<b>No. of preparations* (10 µl QIAEX II per prep)</b>	<b>150</b>	<b>500</b>
QIAEX II Suspension	3 x 0.5 ml	10 x 0.5 ml
Buffer QX1 <sup>†</sup> (with pH indicator)	2 x 100 ml	6 x 100 ml
Buffer PE (concentrate)	2 x 20 ml	6 x 20 ml
Handbook	1	1

\* Routine purifications from gel slices >100 mg which contain fragments <100 bp or where the gels are >2% agarose require additional Buffer QX1 to perform the full number of extractions (see Ordering Information, page 22).

<sup>†</sup> Buffer QX1 contains chaotropic salt which is an irritant. Take appropriate laboratory safety precautions and wear gloves when handling.

## Storage

The QIAEX II Gel Extraction Kit should be stored dry at room temperature (15–25°C). When stored under these conditions and handled appropriately, the kit can be stored for at least 12 months without showing any reduction in performance, capacity, or quality of separation.

## Product Use Limitations

The QIAEX II Gel Extraction Kit is intended for molecular biology applications. This product is neither intended for the diagnosis, prevention, or treatment of a disease, nor has it been validated for such use either alone or in combination with other products. Therefore, the performance characteristics of the product for clinical use (i.e., diagnostic, prognostic, therapeutic, or blood banking) are unknown.

## Product Warranty and Satisfaction Guarantee

QIAGEN guarantees the performance of all products in the manner described in our product literature. The purchaser must determine the suitability of the product for its particular use. Should any product fail to perform satisfactorily due to any reason other than misuse, QIAGEN will replace it free of charge or refund the purchase price. We reserve the right to change, alter, or modify any product to enhance its performance and design. If a QIAGEN product does not meet your expectations, simply call your local Technical Service Department or distributor. We will credit your account or exchange the product — as you wish. Separate conditions apply to QIAGEN scientific instruments, service products, and to products shipped on dry ice. Please inquire for more information.

A copy of QIAGEN terms and conditions can be obtained on request, and is also provided on the back of our invoices. If you have questions about product specifications or performance, please call QIAGEN Technical Services or your local distributor (see back cover or visit [www.qiagen.com](http://www.qiagen.com)).

## Technical Assistance

At QIAGEN, we pride ourselves on the quality and availability of our technical support. Our Technical Service Departments are staffed by experienced scientists with extensive practical and theoretical expertise in sample and assay technologies and the use of QIAGEN products. If you have any questions or experience any difficulties regarding the QIAEX II Gel Extraction Kit or QIAGEN products in general, please do not hesitate to contact us.

QIAGEN customers are also a major source of information regarding advanced or specialized uses of our products. This information is helpful to other scientists as well as to the researchers at QIAGEN. We therefore also encourage you to contact us if you have any suggestions about product performance or new applications and techniques.

For technical assistance and more information, please see our Technical Support Center at [www.qiagen.com/Support](http://www.qiagen.com/Support) or call one of the QIAGEN Technical Service Departments or local distributors (see back cover or visit [www.qiagen.com](http://www.qiagen.com)).

## Safety Information

When working with chemicals, always wear a suitable lab coat, disposable gloves, and protective goggles. For more information, please consult the appropriate material safety data sheets (MSDSs). These are available online in convenient and compact PDF format at [www.qiagen.com/support/MSDS.aspx](http://www.qiagen.com/support/MSDS.aspx) where you can find, view, and print the MSDS for each QIAGEN kit and kit component.

### 24-hour emergency information

Emergency medical information in English, French, and German can be obtained 24 hours a day from:

Poison Information Center Mainz, Germany

Tel: +49-6131-19240

## Product Specifications

Maximum binding capacity: 5 µg DNA per 10 µl QIAEX II suspension

Recovery: 60–95% of DNA fragments 40 bp – 50 kb

Minimum elution volume: 20 µl

## Quality Control

As part of the stringent QIAGEN quality assurance program, the performance of the QIAEX II Gel Extraction Kit is monitored routinely and on a lot-to-lot basis. The QIAEX II Gel Extraction Kit is tested by isolation of DNA fragments of various sizes from either aqueous solution or agarose gel. The quality of the isolated DNA is checked by several assays commonly used for nucleic acids. The DNA binding capacity of QIAEX II resin is tested by determining the recovery from a specific amount of loaded DNA.

## Introduction

The QIAEX II Gel Extraction Kit is designed to extract and purify DNA from any agarose gel in either TAE (Tris-acetate/EDTA) or TBE (Tris-borate/EDTA) buffer without phenol extraction or ethanol precipitation. The QIAEX II silica particles have been optimized to enhance recovery of very small and very large DNA fragments. DNA molecules of 40 bp to 50 kb are adsorbed to QIAEX II particles in the presence of high salt. All non-nucleic acid impurities such as agarose, proteins, salts, and ethidium bromide are removed during washing steps. The pure DNA is efficiently eluted in just 20 µl of Tris buffer or water and is suitable for most subsequent applications, e.g., restriction digestion, labeling, ligation, PCR, sequencing, and microinjection (see Applications using QIAEX II Purified Fragments, page 11).

## Principle

With the QIAEX II Gel Extraction Kit, extraction and purification of DNA fragments are based on solubilization of agarose and selective, quantitative adsorption of nucleic acids to the QIAEX II silica-gel particles in the presence of high salt. Elution of the DNA is accomplished with a low-salt solution such as Tris buffer or water.

### Solubilization of agarose without NaI

The optimized Buffer QX1 in the QIAEX II Gel Extraction Kits efficiently solubilizes agarose, and does not contain NaI. Residual NaI may be difficult to remove from DNA samples, and reduces the efficiency of subsequent enzymatic reactions such as blunt-end ligation.

The standard QIAEX II Gel Extraction protocol is used to extract DNA from 0.3–2% standard or low-melt agarose gels in TAE or TBE buffer. The concentration of agarose in the gel determines the size range of DNA molecules which can be resolved (Table 1).

**Table 1. Separation range in TAE gels containing different concentrations of agarose (1)**

Agarose	Size of DNA fragments separated
0.3%	5 – 60 kb
0.5%	1 – 30 kb
0.7%	0.8 – 12 kb
1.0%	0.5 – 10 kb
1.2%	0.4 – 7 kb
1.5%	0.2 – 3 kb
2.0%	0.05 – 2 kb


A typical agarose gel slice is solubilized by adding 3 volumes of Buffer QX1 to 1 volume of gel (e.g., 300  $\mu$ l of Buffer QX1 is added to 100 mg gel slice) and incubating at 50°C for 10 minutes. The high concentration of a chaotropic salt in Buffer QX1 disrupts hydrogen bonding between sugars in the agarose polymer, allowing solubilization of the gel slice. In addition, the high salt concentration dissociates DNA binding proteins from the DNA fragments.

A  $\leq 2\%$  agarose gel slice is normally solubilized within 2–3 minutes with Buffer QX1 at 50°C. The incubation with QIAEX II is extended to 10 minutes to complete the adsorption of DNA to the QIAEX II particles.

### Efficient DNA adsorption to QIAEX II — salt and pH dependence

Adsorption of DNA to glass, silica gel, or diatomaceous earth in high salt is a well known phenomenon (2). Incubation of a DNA solution in a highly electrolytic environment with large anions (e.g., from chaotropic salts) causes a modification in the structure of water (3), forcing the DNA to adsorb to the silica particles. Adsorption of fragments smaller than 100 bp is enhanced by increasing the salt concentration, while fragments larger than 4 kb are adsorbed at lower salt concentrations.

Adsorption of DNA to silica also depends on pH. Adsorption efficiency is typically 95% if the pH is  $\leq 7.5$ , and is drastically reduced at higher pH (Figure 1). If the pH of the binding mixture is  $>7.5$ , the optimal pH for DNA binding can be achieved by adding a small volume of 3 M sodium acetate, pH 5.0.



**Figure 1. pH dependence of DNA adsorption to silica.** One microgram of a 2.9 kb fragment was adsorbed at different pH values and eluted with 10 mM Tris·Cl, pH 8.5. The graph shows percentage DNA recovery, reflecting the relative adsorption efficiency, versus pH of adsorption.



## QIAEX II Gel Extraction Procedure

Solubilized gel slice  
+ QIAEX II particles



Bind



Collect



Wash



Elute



Pure DNA Fragment

## pH indicator in solubilization and binding Buffer QX1

Buffer QX1 in the QIAEX II Gel Extraction Kit is used for solubilization of agarose gel slices and binding of DNA to the QIAEX II silica particles. Buffer QX1 contains a pH indicator, allowing easy determination of the optimal pH for DNA binding. DNA adsorption requires a pH  $\leq 7.5$ , and the pH indicator appears yellow in this range. If the pH is  $>7.5$ , which can occur if the agarose gel electrophoresis buffer is frequently used or incorrectly prepared, the binding mixture turns orange or violet (Figure 2). This means that the pH of the sample exceeds the buffering capacity of Buffer QX1 and DNA adsorption will be inefficient. In this case, the pH of the binding mixture can easily be corrected by addition of a small volume of 3 M sodium acetate, pH 5.0, before proceeding with the protocol.

The color of the binding mixture allows easy visualization of any unsolubilized agarose, ensuring complete solubilization and maximum yields. The indicator dye does not interfere with DNA binding and is completely removed during the cleanup procedure.



Figure 2. Indicator dye in solubilization and binding Buffer QX1 identifies optimal pH for DNA binding.

## Washing

DNA molecules bind to the QIAEX II particles during the adsorption step, and all non-nucleic acid impurities such as agarose, proteins, ethidium bromide, and salts remain in the supernatant. A high salt wash with Buffer QX1 removes residual agarose, and two washes with ethanol-containing Buffer PE efficiently remove salt contaminants. All traces of supernatant must be carefully removed at each step to eliminate impurities and reduce buffer carryover.

Washed QIAEX II particles carrying adsorbed DNA are pelleted and air-dried at room temperature for 10–15 minutes. Drying the pellet is necessary to remove all traces of residual ethanol which may interfere with subsequent enzymatic reactions.

## Elution in low-salt solutions

Elution efficiency depends on pH and salt concentration. Contrary to adsorption, elution is best under basic conditions and low salt concentrations when using QIAEX II. DNA is typically eluted with 20  $\mu$ l of 10 mM Tris-Cl, pH 8.5, or with water. The maximum elution efficiency is achieved between pH 7.0 and 8.5. When using water to elute, make sure that the pH value is within this range. In addition, DNA must be stored at  $-20^{\circ}\text{C}$  when eluted with water, since DNA may degrade in the absence of a buffering agent. The purified DNA can also be eluted in TE (10 mM Tris-Cl, 1 mM EDTA, pH 8.0), but the EDTA may inhibit subsequent enzymatic reactions.

Elution efficiency also depends on temperature. DNA fragments smaller than 4 kb are recovered with 70–95% efficiency when eluted at room temperature with a 5-min incubation. For efficient elution of DNA fragments larger than 4 kb, the incubation temperature should be increased to  $50^{\circ}\text{C}$ .

After elution, QIAEX II is pelleted by centrifugation. The supernatant containing pure DNA is easily removed from the QIAEX II pellet, ready for use in subsequent applications.

## Applications using QIAEX II purified fragments

DNA purified with QIAEX II is suitable for various applications, for example, restriction digestion, labeling, ligation, transformation, PCR, sequencing, in vitro transcription, and microinjection. Some applications, such as sequencing, blunt-end ligation, in vitro transcription, or microinjection, are highly sensitive to resin carryover and residual salts, and therefore all traces of supernatant must be carefully removed at each wash step. After elution, the supernatant containing the pure DNA should be carefully removed from the QIAEX II pellet. If necessary, the supernatant may be centrifuged a second time to ensure complete removal of all QIAEX II particles.

For sensitive downstream applications such as fluorescent sequencing, QIAquick™ Gel Extraction Kits (see Ordering Information, page 22) are recommended. QIAquick Gel Extraction Kits combine a convenient microspin format with a specially designed silica membrane. The quality of the purified DNA is less dependent on the handling procedures.

# Protocol: Extraction of 40 bp to 50 kb DNA fragments from Agarose Gels

This protocol is designed for the extraction of 40 bp to 50 kb DNA fragments from 0.3–2% standard or low-melt agarose gels in TAE or TBE buffers.

## Important points before starting

- The yellow color of Buffer QX1 indicates a pH  $\leq 7.5$ .
- Add ethanol (96–100%) to Buffer PE before use (see bottle label for volume).
- A heating block or water bath at 50°C is required.
- 3 M sodium acetate, pH 5.0, may be necessary.
- All centrifugation steps are at maximum speed ( $\geq 10,000 \times g$ ,  $\sim 13,000$  rpm) in a conventional, table-top microcentrifuge.
- For DNA fragments larger than 10 kb, mix by gently flicking the tube to avoid shearing the DNA. Do not vortex the tube.

## Procedure

- Excise the DNA band from the agarose gel with a clean, sharp scalpel.**  
Minimize the size of the gel slice by removing excess agarose. Use a 1.5 ml microfuge tube for processing up to 250 mg agarose.
- Weigh the gel slice in a colorless tube. Add 3 volumes of Buffer QX1 to 1 volume of gel for DNA fragments 100 bp – 4 kb; otherwise, follow the table below.**  
For example, add 300  $\mu$ l of Buffer QX1 to each 100 mg of gel.

Table 2. Addition of Buffer QX1

DNA fragments <100 bp	Add 6 volumes of Buffer QX1
DNA fragments >4 kb	Add 3 volumes of Buffer QX1 plus 2 volumes of H <sub>2</sub> O
>2% or Metaphor agarose gels	Add 6 volumes of Buffer QX1

3. **Resuspend QIAEX II by vortexing for 30 s. Add QIAEX II to the sample according to the table below and mix.**

**Table 3. Addition of QIAEX II**

≤2 µg DNA	Add 10 µl of QIAEX II
2–10 µg DNA	Add 30 µl of QIAEX II
Each additional 10 µg DNA	Add additional 30 µl of QIAEX II

4. **Incubate at 50°C for 10 min to solubilize the agarose and bind the DNA. Mix by vortexing\* every 2 min to keep QIAEX II in suspension. Check that the color of the mixture is yellow.**

If the color of the mixture is orange or purple, add 10 µl 3 M sodium acetate, pH 5.0, and mix. The color should turn to yellow. The incubation should then be continued for an additional 5 min at least.

The adsorption of DNA to QIAEX II particles is only efficient at pH ≤7.5. Buffer QX1 now contains a pH indicator which is yellow at pH ≤7.5, and orange or violet at higher pH, allowing easy determination of the optimal pH for DNA binding.

5. **Centrifuge the sample for 30 s and carefully remove supernatant with a pipet.**
6. **Wash the pellet with 500 µl of Buffer QX1.**

Resuspend the pellet by vortexing\*. Centrifuge the sample for 30 s and remove all traces of supernatant with a pipet. This wash step removes residual agarose contaminants.

7. **Wash the pellet twice with 500 µl of Buffer PE.**

Resuspend the pellet by vortexing\*. Centrifuge the sample for 30 s and carefully remove all traces of supernatant with a pipet. These washing steps remove residual salt contaminants.

8. **Air-dry the pellet for 10–15 min or until the pellet becomes white.**

If 30 µl of QIAEX II suspension is used, air-dry the pellet for approximately 30 min. Do not vacuum dry, as this may cause overdrying. Overdrying the QIAEX II pellet may result in decreased elution efficiency.

9. **To elute DNA, add 20 µl of 10 mM Tris-Cl, pH 8.5 or H<sub>2</sub>O and resuspend the pellet by vortexing\*. Incubate according to the table in the next page.**

\* For fragments larger than 10 kb, resuspend the pellet by inverting and flicking the tube. Vortexing can cause shearing of large DNA fragments.

**Table 4. Incubation conditions**

DNA fragments $\leq 4$ kb	Incubate at room temp. for 5 min
DNA fragments 4–10 kb	Incubate at 50°C for 5 min
DNA fragments >10 kb	Incubate at 50°C for 10 min

Elution efficiency is dependent on pH. The maximum elution efficiency is achieved between pH 7.0 and 8.5. When using water for elution, make sure that the pH is within this range, and store DNA at  $-20^{\circ}\text{C}$  as DNA may degrade in the absence of a buffering agent. The purified DNA can also be eluted in TE buffer (10 mM Tris-Cl, 1 mM EDTA, pH 8.0) but the EDTA may inhibit subsequent enzymatic reactions.

**10. Centrifuge for 30 s. Carefully pipet the supernatant into a clean tube.**

The supernatant now contains the purified DNA.

**11. Optional: repeat steps 9 and 10 and combine the eluates.**

A second elution step will increase the yield by approximately 10–15%.

# Protocol: Extraction of DNA fragments from Polyacrylamide Gels

## Important points before starting:

- The yellow color of Buffer QX1 indicates a pH  $\leq 7.5$ .
- Add ethanol (96–100%) to Buffer PE before use (see bottle label for volume).
- A heating block or water bath at 50°C is required.
- 3 M sodium acetate, pH 5.0, may be necessary.
- All centrifugation steps are at maximum speed ( $\geq 10,000 \times g$ ,  $\sim 13,000$  rpm) in a conventional, table-top microcentrifuge.
- Prepare diffusion buffer: 0.5 M ammonium acetate; 10 mM magnesium acetate; 1 mM EDTA, pH 8.0; 0.1% SDS.
- A disposable plastic column or a syringe barrel containing either a Whatman® GF/C filter or packed, siliconized glass wool will be required for each extraction.

## Procedure

1. **Excise the gel slice containing the DNA band with a clean, sharp scalpel.**  
Minimize the size of the gel slice by removing excess polyacrylamide.
2. **Weigh the gel slice. Add 1–2 volumes of diffusion buffer to 1 volume of gel (i.e., 100–200  $\mu$ l for each 100 mg of gel).**
3. **Incubate at 50°C for 30 min.**
4. **Microcentrifuge the sample for 1 min.**
5. **Carefully remove the supernatant using a pipet or a drawn-out Pasteur pipet. Pass the supernatant through a disposable plastic column or a syringe containing either a Whatman GF/C filter or packed, siliconized glass wool to remove any residual polyacrylamide.**
6. **Calculate the approximate volume of the recovered supernatant.**
7. **For DNA fragments <100 bp, add 6 volumes of Buffer QX1 to 1 volume of sample. Otherwise, add 3 volumes Buffer QX1. Check that the color of the mixture is yellow.**

If the color of the mixture is orange or purple, add 10  $\mu$ l 3 M sodium acetate, pH 5.0, and mix. The color should turn to yellow.

The adsorption of DNA to QIAEX II particles is only efficient at pH  $\leq 7.5$ . Buffer QX1 now contains a pH indicator which is yellow at pH  $\leq 7.5$ , and orange or violet at higher pH, allowing easy determination of optimal pH for DNA-binding.

8. **Resuspend QIAEX II by vortexing for 30 s.**

9. Add 10  $\mu$ l of QIAEX II and mix. Incubate at room temperature for 10 min. Vortex every 2 min to keep QIAEX II in suspension.
10. Centrifuge the sample for 30 s and remove supernatant.
11. Wash the pellet twice with 500  $\mu$ l of Buffer PE.
12. Air-dry the pellet for 10–15 min or until the pellet becomes white.

Do not vacuum dry, as this may cause overdrying. Overdrying the QIAEX II pellet may result in decreased elution efficiency.

13. To elute DNA, add 20  $\mu$ l of 10 mM Tris-Cl, pH 8.5, or H<sub>2</sub>O and resuspend the pellet by vortexing. Incubate for 5 min at room temperature.

Elution efficiency is dependent on pH. The maximum elution efficiency is achieved between pH 7.0 and 8.5. When using water for elution, make sure that the pH is within this range, and store DNA at –20°C as DNA may degrade in the absence of a buffering agent. The purified DNA can also be eluted in TE buffer (10 mM Tris-Cl, 1 mM EDTA, pH 8.0) but the EDTA may inhibit subsequent enzymatic reactions.

14. Centrifuge for 30 s. Carefully transfer the supernatant into a clean tube.

The supernatant now contains pure DNA.

15. Optional: repeat steps 13 and 14 and combine the eluates.

A second elution step will increase the yield by approximately 10–15%.



## Protocol: Desalting and Concentrating DNA Solutions

QIAEX II Gel Extraction Kit can be used to purify and concentrate DNA fragments from 40 bp to 50 kb from aqueous solutions without phenol extraction or ethanol precipitation. Enzymes such as phosphatases, restriction endonucleases, polymerases, as well as dNTPs, and salts are removed. For direct purification from PCR reactions, use the QIAquick PCR Purification Kit (see Ordering Information, page 22) to recover PCR products and efficiently remove primers.

### Important points before starting

- The yellow color of Buffer QX1 indicates a pH  $\leq 7.5$ .
- Add ethanol (96–100%) to Buffer PE before use (see bottle label for volume).
- All centrifugation steps are at maximum speed ( $\geq 10,000 \times g$ ,  $\sim 13,000$  rpm) in a conventional table-top microcentrifuge.
- For DNA fragments larger than 10 kb, mix by gently flicking the tube to avoid shearing the DNA. Do not vortex the tube.

### Procedure

1. **Transfer the sample to a colorless tube. Add 3 volumes of Buffer QX1 to 1 volume of sample.**

For DNA fragments 100 bp – 4 kb; otherwise follow the table below.

**Table 5. Addition of Buffer QX1**

DNA fragments <100 bp	Add 6 volumes of Buffer QX1
DNA fragments >4 kb	Add 3 volumes of Buffer QX1 plus 2 volumes of H <sub>2</sub> O

2. **Check that the color of the sample mixture is yellow.**

If the color of the mixture is orange or purple, add 10  $\mu$ l 3 M sodium acetate, pH 5.0, and mix. The color should now be yellow.

The adsorption of DNA to QIAEX II particles is only efficient at pH  $\leq 7.5$ . Buffer QX1 now contains a pH indicator which is yellow at pH  $\leq 7.5$ , and orange or violet at higher pH, allowing easy determination of the optimal pH for DNA-binding.

3. **Resuspend QIAEX II by vortexing 30 s.**

4. Add 10  $\mu$ l of QIAEX II per 5  $\mu$ g of DNA and mix\*. Incubate at room temperature for 10 min. Mix every 2 min to keep QIAEX II in suspension.
5. Centrifuge the sample for 30 s and remove supernatant.
6. Wash the pellet twice with 500  $\mu$ l of Buffer PE.
7. Air-dry the pellet for 10–15 min or until the pellet becomes white.  
Do not vacuum dry, as this may cause overdrying. Overdrying the QIAEX II pellet may result in decreased elution efficiency.
8. To elute DNA, add 20  $\mu$ l of 10 mM Tris-Cl, pH 8.5, or H<sub>2</sub>O and resuspend the pellet by vortexing\*. Incubate according to the table below.

**Table 6. Incubation conditions**

DNA fragments $\leq$ 4 kb	Incubate at room temp. for 5 min
DNA fragments 4–10 kb	Incubate at 50°C for 5 min
DNA fragments >10 kb	Incubate at 50°C for 10 min

Elution efficiency is dependent on pH. The maximum elution efficiency is achieved between pH 7.0 and 8.5. When using water for elution, make sure that the pH is within this range, and store DNA at  $-20^{\circ}\text{C}$  as DNA may degrade in the absence of a buffering agent. The purified DNA can also be eluted in TE buffer (10 mM Tris-Cl, 1 mM EDTA, pH 8.0) but the EDTA may inhibit subsequent enzymatic reactions.

9. **Centrifuge for 30 s. Carefully transfer the supernatant into a clean tube.**  
The supernatant now contains pure DNA.
10. **Optional: repeat steps 8 and 9 and combine the eluates.**  
A second elution step will increase the yield by approximately 10–15%.

\* For fragments larger than 10 kb, resuspend the pellet by inverting and flicking the tube. Vortexing can cause shearing of large DNA fragments.

# Troubleshooting Guide

This troubleshooting guide may be helpful in solving any problems that may arise. For more information, see also the Frequently Asked Questions page at our Technical Support Center: [www.qiagen.com/FAQ/FAQList.aspx](http://www.qiagen.com/FAQ/FAQList.aspx). The scientists in QIAGEN Technical Services are always happy to answer any questions you may have about either the information and protocols in this handbook or sample and assay technologies (for contact information, see back cover or visit [www.qiagen.com](http://www.qiagen.com)).

---

## Comments and suggestions

---

### Poor DNA recovery

- |   |  |
|---|--|
| a) Ethanol was not added to concentrated Buffer PE before use                       | Repeat procedure with correctly prepared Buffer PE.  |
| b) pH of electrophoresis buffer too high (binding mixture turns orange or violett). | The electrophoresis buffer has been repeatedly used or incorrectly prepared, resulting in a sample pH that exceeds the buffering capacity of Buffer QX1 and leads to inefficient DNA binding. Add 10 µl of 3 M sodium acetate, pH 5.0, to the sample and mix. The color of the mixture should turn yellow indicating the correct pH for DNA binding. For binding mixtures with only small color changes (slight orange color), add the sodium acetate. |
| c) Inappropriate elution buffer   | DNA will only be eluted in the presence of low salt buffer (e.g., 10 mM Tris-Cl, pH 8.5) or water. Check the pH and salt concentration of the elution buffer.  |
| d) Incomplete solubilization of gel slice in Buffer QX1                             | Mix the tube every 2 min during the solubilization step. Large gel slices may require a few extra minutes at 50°C to solubilize. DNA will remain in any undissolved agarose.   |
| e) Low salt concentration during adsorption   | Sufficient salt must be present during the adsorption step. Weigh the gel slice weight accurately and add the appropriate amount of Buffer QX1.  |

## Comments and suggestions

- |    |   |  |
|----|---|--|
| f) | Insufficient amount of QIAEX II                                 | Enough QIAEX II must be present at the adsorption step to bind DNA. See Table 3 on page 13 for details.  |
| g) | Buffer QX1 incompletely removed before wash step with Buffer PE | If Buffer QX1 is not completely removed, the subsequent Buffer PE washing steps may not efficiently remove all traces of salt. The presence of high concentrations of salt may prevent complete elution of DNA from the QIAEX II particles. Additionally, residual salt may result in poor performance of the DNA in subsequent enzymatic reactions. |
| h) | Over-dried QIAEX II pellet                                      | Do not use a vacuum dryer to dry the QIAEX II pellet. If pellet is dried too extensively, the DNA may irreversibly bind to the QIAEX II particles, resulting in decreased recovery.  |
| i) | Insufficient volume of elution buffer                           | Increase volume of elution buffer in proportion to amount of extra QIAEX II added. An optional extra elution step will increase yield by approximately 10–15%.   |
| j) | Incorrect volume of Buffer QX1 added for solubilization         | See Table 2 on page 12 for details.  |

### DNA does not perform well in subsequent reactions

- |    |  |   |
|----|--|---|
| a) | Incomplete removal of Buffer QX1 supernatant           | Salt from Buffer QX1 may be carried over and inhibit subsequent enzymatic reactions. Wash twice with Buffer PE.   |
| b) | Incomplete removal of Buffer PE supernatant            | Residual ethanol may reduce elution efficiency. Carefully remove all Buffer PE to enable pellet to dry sufficiently before elution.   |
| c) | Incomplete removal of QIAEX II particles after elution | QIAEX II particles may bind proteins under low salt conditions, and reduce enzyme activity in subsequent reactions. Centrifuge again and take the supernatant to remove all traces of QIAEX II. |

### DNA contains residual ethanol

#### (samples float out of wells of agarose gels)

- |    |                                 |  |
|----|---------------------------------|--|
| a) | QIAEX II pellet was under-dried | Buffer PE was not completely removed by drying. Dry the DNA eluate briefly in a vacuum dryer to remove residual ethanol. |
|----|---------------------------------|--|

### DNA appears to “smear” on an agarose gel.

- |                                    |  |
|------------------------------------|--|
| a) Sample well is over-loaded      | Smiling and smearing of DNA bands may occur when too much sample is loaded in one well. Use less DNA per well or use a larger well.  |
| b) Vigorous handling of DNA        | Vortexing and vigorous pipetting of fragments above 10 kb should be avoided throughout. QIAEX II can be easily resuspended by flicking the microfuge tube. Very large DNA fragments can be protected from shearing by keeping the QIAEX II pellet intact during the washing steps. The wash buffers should remain on the QIAEX II pellet for 5 min to allow complete diffusion.  |
| c) Eluate contains denatured ssDNA | Denatured ssDNA appears as a smaller smeared band on an analytical gel. Use the eluted DNA to prepare the subsequent enzymatic reaction but omit the enzyme. To reanneal the ssDNA, incubate the reaction mixture at 95°C for 2 min, and allow the tube to cool slowly to room temperature. Add the enzyme and proceed as usual. Alternatively, the DNA can be eluted in 10 mM Tris buffer containing 10 mM NaCl. The salt and buffering agent promote the renaturation of DNA strands. However, the salt concentration of the eluate must then be considered for subsequent applications. |

## References

1. Sambrook, J., Fritsch, E.F., and Maniatis, T. (1989) *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor, NY: Cold Spring Harbor Laboratory Press.
2. Vogelstein, B., and Gillespie, D. (1979) Preparative and analytical purification of DNA from agarose. *Proc. Natl. Acad. Sci. USA* **76**, 615–619.
3. Hamaguchi, K., and Geiduschek, E.P. (1962) The effect of electrolytes on the stability of deoxyribonucleate helix. *J. Am. Chem. Soc.* **84**, 1329–1337.

# Ordering Information

Product	Contents	Cat. no.
QIAEX II Gel Extraction Kit (150)	For 150 extractions*: 3 x 0.5 ml QIAEX II Suspension, Buffers	20021
QIAEX II Gel Extraction Kit (500)	For 500 extractions*: 5 x 1.0 ml QIAEX II Suspension, Buffers	20051
QIAEX II Suspension	3 x 0.5 ml QIAEX II Suspension	20902
Buffer QX1	500 ml Solubilization and Binding Buffer (with pH indicator)	20912
Buffer PE	100 ml Wash Buffer (concentrate; final volume 500 ml)	19065
<b>Related products</b>		
QIAquick PCR Purification Kit (50) <sup>†</sup>	50 QIAquick Spin Columns, Buffers, Collection Tubes (2 ml)	28104
QIAquick Gel Extraction Kit (50) <sup>†</sup>	50 QIAquick Spin Columns, Buffers, Collection Tubes (2 ml)	28704
QIAquick Nucleotide Removal Kit (50) <sup>†</sup>	50 QIAquick Spin Columns, Buffers, Collection Tubes (2 ml)	28304
MinElute PCR Purification Kit (50) <sup>†</sup>	50 MinElute Spin Columns, Buffers, Collection Tubes (2 ml)	28004
MinElute Gel Extraction Kit (50) <sup>†</sup>	50 MinElute Spin Columns, Buffers, Collection Tubes (2 ml)	28604
MinElute Reaction Cleanup Kit (50) <sup>†</sup>	50 MinElute Spin Columns, Buffers, Collection Tubes (2 ml)	28204

All kits are intended for molecular biology applications. These products are neither intended for the diagnosis, prevention, or treatment of a disease, nor have they been validated for such use either alone or in combination with other products.

\* Routine purifications from gel slices >100 mg which contain fragments <100 bp or where the gels contain >2% agarose require additional Buffer QX1 to perform the full number of extractions.

<sup>†</sup> Larger kit sizes/formats available; see [www.qiagen.com](http://www.qiagen.com).

Trademarks: QIAGEN®, QIAEX®, QIAquick®, MinElute® (QIAGEN Group); Whatman® (Whatman International Ltd.).

Registered names, trademarks, etc. used in this document, even when not specifically marked as such, are not to be considered unprotected by law.

Purchase of this product is not accompanied by a license under patents owned by Roche Molecular Systems, Inc. and F. Hoffmann-La Roche Ltd. No rights to real-time PCR or to any primers, probes or pathogens are conveyed expressly, by implication or by estoppel.

### **Limited License Agreement**

Use of this product signifies the agreement of any purchaser or user of the QIAEX II Gel Extraction Kit to the following terms:

1. The QIAEX II Gel Extraction Kit may be used solely in accordance with the *QIAEX II Handbook* and for use with components contained in the Kit only. QIAGEN grants no license under any of its intellectual property to use or incorporate the enclosed components of this Kit with any components not included within this Kit except as described in the *QIAEX II Handbook* and additional protocols available at [www.qiagen.com](http://www.qiagen.com).
2. Other than expressly stated licenses, QIAGEN makes no warranty that this Kit and/or its use(s) do not infringe the rights of third-parties.
3. This Kit and its components are licensed for one-time use and may not be reused, refurbished, or resold.
4. QIAGEN specifically disclaims any other licenses, expressed or implied other than those expressly stated.
5. The purchaser and user of the Kit agree not to take or permit anyone else to take any steps that could lead to or facilitate any acts prohibited above. QIAGEN may enforce the prohibitions of this Limited License Agreement in any Court, and shall recover all its investigative and Court costs, including attorney fees, in any action to enforce this Limited License Agreement or any of its intellectual property rights relating to the Kit and/or its components.

For updated license terms, see [www.qiagen.com](http://www.qiagen.com).

© 2008 QIAGEN, all rights reserved.

[www.qiagen.com](http://www.qiagen.com)

**Australia** = Orders 03-9840-9800 = Fax 03-9840-9888 = Technical 1-800-243-066

**Austria** = Orders 0800/28-10-10 = Fax 0800/28-10-19 = Technical 0800/28-10-11

**Belgium** = Orders 0800-79612 = Fax 0800-79611 = Technical 0800-79556

**Brazil** = Orders 0800-557779 = Fax 55-11-5079-4001 = Technical 0800-557779

**Canada** = Orders 800-572-9613 = Fax 800-713-5951 = Technical 800-DNA-PREP (800-362-7737)

**China** = Orders 0086-21-3865-3865 = Fax 0086-21-3865-3965 = Technical 800-988-0325, 800-988-0327

**Denmark** = Orders 80-885945 = Fax 80-885944 = Technical 80-885942

**Finland** = Orders 0800-914416 = Fax 0800-914415 = Technical 0800-914413

**France** = Orders 01-60-920-926 = Fax 01-60-920-925 = Technical 01-60-920-930 = Offers 01-60-920-928

**Germany** = Orders 02103-29-12000 = Fax 02103-29-22000 = Technical 02103-29-12400

**Hong Kong** = Orders 800 933 965 = Fax 800 930 439 = Technical 800 930 425

**Ireland** = Orders 1800-555-049 = Fax 1800-555-048 = Technical 1800-555-061

**Italy** = Orders 02-33430-420 = Fax 02-33430-426 = Technical 800-787980

**Japan** = Telephone 03-6890-7300 = Fax 03-5547-0818 = Technical 03-6890-7300

**Korea (South)** = Orders 1544 7145 = Fax 1544 7146 = Technical 1544 7145

**Luxembourg** = Orders 8002-2076 = Fax 8002-2073 = Technical 8002-2067

**Mexico** = Orders 01-800-7742-639 = Fax 01-800-1122-330 = Technical 01-800-7742-639

**The Netherlands** = Orders 0800-0229592 = Fax 0800-0229593 = Technical 0800-0229602

**Norway** = Orders 800-18859 = Fax 800-18817 = Technical 800-18712

**Singapore** = Orders 65-67775366 = Fax 65-67785177 = Technical 65-67775366

**Spain** = Orders 91-630-7050 = Fax 91-630-5145 = Technical 91-630-7050

**Sweden** = Orders 020-790282 = Fax 020-790582 = Technical 020-798328

**Switzerland** = Orders 055-254-22-11 = Fax 055-254-22-13 = Technical 055-254-22-12

**UK** = Orders 01293-422-911 = Fax 01293-422-922 = Technical 01293-422-999

**USA** = Orders 800-426-8157 = Fax 800-718-2056 = Technical 800-DNA-PREP (800-362-7737)

